# Ku70 Rabbit pAb

**Catalog NO.: A0883** **11 Publications**

## Basic Information

### Observed MW
70kDa

### Calculated MW
70kDa

### Category
Polyclonal Antibodies

### Applications
WB, IHC-P, IF/ICC, ELISA

### Cross-Reactivity
Human, Mouse, Rat

### Conjugation
Unconjugated

## Background

The p70/p80 autoantigen is a nuclear complex consisting of two subunits with molecular masses of approximately 70 and 80 kDa. The complex functions as a single-stranded DNA-dependent ATP-dependent helicase. The complex may be involved in the repair of nonhomologous DNA ends such as that required for double-strand break repair, transposition, and V(D)J recombination. High levels of autoantibodies to p70 and p80 have been found in some patients with systemic lupus erythematosus.

## Recommended Dilutions

| | |
|---|---|
| **WB** | 1:3000 - 1:18000 |
| **IHC-P** | 1:50 - 1:200 |
| **IF/ICC** | 1:50 - 1:200 |

### ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID
2547

### Swiss Prot
P12956

### Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms
ML8; KU70; TLAA; CTC75; CTCBF; G22P1; Ku70

## Contact

## Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com


**Source**
Rabbit

**Isotype**
IgG


**Purification**
Affinity purification

**Storage**
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## Validation Data






Western blot analysis of various lysates using Ku70 Rabbit pAb (A0883) at 1:3000 dilution incubated at room temperature for 1.5 hours.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.




Western blot analysis of lysates from C6 cells using Ku70 Rabbit pAb (A0883) at 1:3000 dilution incubated at room temperature for 1.5 hours.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 45s.








Immunohistochemistry analysis of paraffin-embedded Rat brain using Ku70 Rabbit pAb (A0883) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.


Immunohistochemistry analysis of paraffin-embedded Human breast using Ku70 Rabbit pAb (A0883) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Human leiomyoma of uterus using Ku70 Rabbit pAb (A0883) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

## Validation Data



Immunohistochemistry analysis of paraffin-embedded Mouse lung using Ku70 Rabbit pAb (A0883) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Confocal immunofluorescence analysis of U2OS cells using Ku70 Rabbit pAb (A0883) at dilution of 1:100. Blue: DAPI for nuclear staining.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.