

# NARS Rabbit pAb

Catalog NO.: A11668

## Basic Information

### Observed MW

63kDa

### Calculated MW

63kDa

### Category

Polyclonal Antibodies

### Applications

WB, ELISA

### Cross-Reactivity

Human, Mouse

### Conjugation

Unconjugated

## Background

Aminoacyl-tRNA synthetases are a class of enzymes that charge tRNAs with their cognate amino acids. Asparaginyl-tRNA synthetase is localized to the cytoplasm and belongs to the class II family of tRNA synthetases. The N-terminal domain represents the signature sequence for the eukaryotic asparaginyl-tRNA synthetases.

## Recommended Dilutions

**WB** 1:500 - 1:2000

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

4677

### Swiss Prot

O43776

### Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms

NARS; ASNRS; NEDMILG; NEDMILEG


## Contact

## Product Information

|                                     |  |                      |
|-------------------------------------|--|----------------------|
| <input checked="" type="checkbox"/> |  | order@abclonal.com   |
| <input checked="" type="checkbox"/> |  | service@abclonal.com |
| <input checked="" type="checkbox"/> |  | www.abclonal.com     |

|                                                       |                |                       |
|-------------------------------------------------------|----------------|-----------------------|
| <b>Source</b>                                         | <b>Isotype</b> | <b>Purification</b>   |
| Rabbit                                                | IgG            | Affinity purification |
| <b>Storage</b>                                        |                |                       |
| Store at -20°C. Avoid freeze / thaw cycles.           |                |                       |
| Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. |                |                       |

Validation Data



Western blot analysis of various lysates using NARS Rabbit pAb (A11668) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 3s.

