

NCKAP1 Rabbit pAb

Catalog NO.: A12229

Basic Information

Observed MW

125kDa

Calculated MW

129kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Background

Contributes to small GTPase binding activity. Involved in Rac protein signal transduction; positive regulation of Arp2/3 complex-mediated actin nucleation; and positive regulation of lamellipodium assembly. Located in extracellular exosome and focal adhesion. Part of SCAR complex.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

10787

Swiss Prot

Q9Y2A7

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

HEM2; NAP1; NAP125; p125Nap1; NCKAP1


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.		

Validation Data



Western blot analysis of various lysates using NCKAP1 Rabbit pAb (A12229) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.

