

RYBP Rabbit pAb

Catalog NO.: A14605 **2 Publications**

Basic Information

Observed MW

33kDa

Calculated MW

25kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Mouse, Rat

Conjugation

Unconjugated

Background

Predicted to enable DNA binding activity and transcription coregulator activity. Involved in several processes, including histone H2A monoubiquitination; negative regulation of proteasomal ubiquitin-dependent protein catabolic process; and positive regulation of transcription, DNA-templated. Located in nucleoplasm. Colocalizes with PcG protein complex.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

23429

Swiss Prot

Q8N488

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

AAP1; DEDAF; YEAF1; APAP-1; RYBP

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using RYBP Rabbit pAb (A14605) at 1:1000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 10s.

