

TRIP4 Rabbit pAb

Catalog NO.: A15352

Basic Information

Observed MW

66kDa

Calculated MW

66kDa

Category

Polyclonal Antibodies

Applications

WB, IHC-P, IP, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Background

This gene encodes a subunit of the tetrameric nuclear activating signal cointegrator 1 (ASC-1) complex, which associates with transcriptional coactivators, nuclear receptors and basal transcription factors to facilitate nuclear receptors-mediated transcription. This protein is localized in the nucleus and contains an E1A-type zinc finger domain, which mediates interaction with transcriptional coactivators and ligand-bound nuclear receptors, such as thyroid hormone receptor and retinoid X receptor alpha, but not glucocorticoid receptor. Mutations in this gene are associated with spinal muscular atrophy with congenital bone fractures-1 (SMABF1).

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

IP

0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Contact

Immunogen Information

Gene ID	Swiss Prot
9325	Q15650

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms


ASC1; ASC-1; MDCDC; SMABF1; ZC2HC5; HsT17391; TRIP4

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.		


Validation Data




Western blot analysis of various lysates using TRIP4 Rabbit pAb (A15352) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.




Immunoprecipitation analysis of 200 µg extracts of A-549 cells using 3 µg TRIP4 antibody (A15352). Western blot was performed from the immunoprecipitate using TRIP4 antibody (A15352) at a dilution of 1:1000.



Immunohistochemistry analysis of paraffin-embedded Rat heart using TRIP4 Rabbit pAb (A15352) at dilution of 1:100 (40x lens). Microwave antigen



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using TRIP4 Rabbit pAb (A15352) at dilution of 1:100 (40x lens).



Immunohistochemistry analysis of paraffin-embedded Human esophageal using TRIP4 Rabbit pAb (A15352) at dilution of 1:100 (40x lens).

retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.