

TRIM62 Rabbit pAb

Catalog NO.: A15855

Basic Information

Observed MW

54kDa

Calculated MW

54kDa

Category

Polyclonal Antibodies

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Background

Enables identical protein binding activity; transcription coactivator activity; and ubiquitin-protein transferase activity. Involved in several processes, including negative regulation of viral transcription; positive regulation of NF-kappaB transcription factor activity; and positive regulation of antifungal innate immune response. Is active in cytoplasm.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

55223

Swiss Prot

Q9BVG3

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

DEAR1; TRIM62

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% sodium azide, 50% glycerol, pH 7.3.

Validation Data



Western blot analysis of various lysates using TRIM62 Rabbit pAb (A15855) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded Human colon using TRIM62 Rabbit pAb (A15855) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

