

XPO7 Rabbit pAb

Catalog NO.: A16108

Basic Information

Observed MW

124kDa

Calculated MW

124kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Background

The transport of protein and large RNAs through the nuclear pore complexes (NPC) is an energy-dependent and regulated process. The import of proteins with a nuclear localization signal (NLS) is accomplished by recognition of one or more clusters of basic amino acids by the importin-alpha/beta complex; see MIM 600685 and MIM 602738. The small GTPase RAN (MIM 601179) plays a key role in NLS-dependent protein import. RAN-binding protein-16 is a member of the importin-beta superfamily of nuclear transport receptors.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

23039

Swiss Prot

Q9UIA9

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

EXP7; RANBP16; XPO7


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.		

Validation Data



Western blot analysis of various lysates using XPO7 Rabbit pAb (A16108) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.

