

# PNLIPRP1 Rabbit pAb

Catalog NO.: A17520

## Basic Information

### Observed MW

52kDa

### Calculated MW

52kDa

### Category

Polyclonal Antibodies

### Applications

WB, IF-P, IHC-P, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### Conjugation

Unconjugated

## Background

Predicted to enable calcium ion binding activity. Predicted to be involved in lipid catabolic process. Predicted to be located in extracellular region. Biomarker of prostate cancer.

## Recommended Dilutions

**WB** 1:500 - 1:2000

**IF-P** 1:50 - 1:200

**IHC-P** 1:100 - 1:200

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

5407

### Swiss Prot

P54315

### Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms

PLRP1; PNLIPRP1


## Contact

## Product Information


|                                     |  |                      |
|-------------------------------------|--|----------------------|
| <input checked="" type="checkbox"/> |  | order@abclonal.com   |
| <input checked="" type="checkbox"/> |  | service@abclonal.com |
| <input checked="" type="checkbox"/> |  | www.abclonal.com     |

|                                                       |                |                       |
|-------------------------------------------------------|----------------|-----------------------|
| <b>Source</b>                                         | <b>Isotype</b> | <b>Purification</b>   |
| Rabbit                                                | IgG            | Affinity purification |
| <b>Storage</b>                                        |                |                       |
| Store at -20°C. Avoid freeze / thaw cycles.           |                |                       |
| Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. |                |                       |


Validation Data




Western blot analysis of various lysates using PNLIPRP1 Rabbit pAb (A17520) at 1:1000 dilution.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 5s.






Immunohistochemistry analysis of paraffin-embedded Rat pancreas using PNLIPRP1 Rabbit pAb (A17520) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human appendix using PNLIPRP1 Rabbit pAb (A17520) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse pancreas using PNLIPRP1 Rabbit pAb (A17520) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of paraffin-embedded rat pancreas using PNLIPRP1 Rabbit pAb (A17520) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of paraffin-embedded rat pancreas using PNLIPRP1 Rabbit pAb (A17520) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of paraffin-embedded mouse pancreas using PNLIPRP1 Rabbit pAb (A17520) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.