

TPST2 Rabbit pAb

Catalog NO.: A18141

Basic Information

Observed MW

42kDa

Calculated MW

42kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Human,Mouse

Conjugation

Unconjugated

Background

The protein encoded by this gene catalyzes the O-sulfation of tyrosine residues within acidic regions of proteins. The encoded protein is a type II integral membrane protein found in the Golgi body. Alternative splicing produces multiple transcript variants encoding distinct isoforms.

Recommended Dilutions

WB

1:100 - 1:500

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

8459

Swiss Prot

O60704

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

TPST-2; TANGO13B; TPST2

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from U-2 OS cells using TPST2 Rabbit pAb (A18141) at 1:1000 dilution incubated overnight at 4°C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020)

Exposure time: 60s.

