

K63-linkage Specific Polyubiquitin Rabbit pAb

Catalog NO.: A18164 **11 Publications**

Basic Information

Observed MW
15-200kDa

Calculated MW

Category
Other Modified Antibodies

Applications
DB, ELISA

Cross-Reactivity
Human, Mouse, Rat, Other
(Wide Range Predicted)

Background

Ubiquitination, onetype of the most common post-translational modification, mediates the regulation of protein homeostasis in vivo. Since ubiquitin itself contains multiple lysine residues and one N-terminal free amino group, eight types of ubiquitin chains can be formed. K63-linked polyubiquitin chains exert nonproteolytic functions in vivo, such as protein trafficking, kinase/phosphatase activation, and DNA damage control, all of which might be important in regulation of cancer survival and development (PMID: 19217402, PMCID: PMC3008764) .

Recommended Dilutions

DB 1:500 - 1:1000

ELISA
Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID **Swiss Prot**

Immunogen
This information is considered to be commercially sensitive.

Synonyms

Contact

Product Information

|

order@abclonal.com

|

service@abclonal.com

|

www.abclonal.com


Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Dot-blot analysis of all sorts of peptides using K63-linkage Specific Polyubiquitin antibody (A18164) at 1:1000 dilution.

