

K11-linkage Specific Polyubiquitin Rabbit pAb

Catalog NO.: A18197 **2 Publications**

Basic Information

Observed MW

Calculated MW

Category

Other Modified Antibodies

Applications

WB, DB, ELISA

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

Background

Polyubiquitination is a posttranslational modification where ubiquitin chains containing isopeptide bonds linking one of seven ubiquitin lysines with the C terminus of an adjoining ubiquitin are covalently attached to proteins. A crystal structure of K11-linked diubiquitin demonstrates a distinct conformation from K48- or K63-linked diubiquitin. It is important that K11-linked ubiquitin chains as critical regulators of mitotic protein degradation.

Recommended Dilutions

WB 1:500 - 1:2000

DB 1:500 - 1:1000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID **Swiss Prot**

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.		

Validation Data



Western blot analysis of various lysates using K11-linkage Specific Polyubiquitin Rabbit pAb (A18197) at 1:500 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25μg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.



Dot-blot analysis of all sorts of peptides using K11-linkage Specific Polyubiquitin antibody (A18197) at 1:1000 dilution.

