

Lambda Light chain Rabbit mAb

Catalog NO.: A19831 **Recombinant**

Basic Information

Observed MW

25kDa

Calculated MW

12kDa

Category

Monoclonal Antibodies

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Clone number

ARC2357

Background

Predicted to be involved in immune response. Located in extracellular exosome.

Recommended Dilutions

WB 1:1000 - 1:6000

IHC-P 1:100 - 1:1000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

28820

Swiss Prot

P01701

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

V1-19; IGLV151; Lambda Light chain

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from Mouse spleen, using Lambda Light chain Rabbit mAb (A19831) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Western blot analysis of lysates from Rat spleen, using Lambda Light chain Rabbit mAb (A19831) at 1:1000 dilution.



Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded Human cervix cancer using Lambda Light chain Rabbit mAb (A19831) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Human tonsil using Lambda Light chain Rabbit mAb (A19831) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.