

SATB2 Rabbit mAb

Catalog NO.: A19837 **Recombinant**

Basic Information

Observed MW

100 kDa

Calculated MW

83 kDa

Category

Monoclonal Antibodies

Applications

WB, IHC-P, IP, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Clone number

ARC2363

Background

This gene encodes a DNA binding protein that specifically binds nuclear matrix attachment regions. The encoded protein is involved in transcription regulation and chromatin remodeling. Defects in this gene are associated with isolated cleft palate and cognitive disability. Alternate splicing results in multiple transcript variants that encode the same protein.

Recommended Dilutions

WB 1:1000 - 1:2000

IHC-P 1:50 - 1:200

IP

0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Contact

Immunogen Information

Gene ID

23314

Swiss Prot

Q9UPW6

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

GLSS; DEL2Q32Q33; C2DELq32q33; SATB2

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from K-562 cells using SATB2 Rabbit mAb (A19837) at 1:1000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Western blot analysis of lysates from Mouse brain using SATB2 Rabbit mAb (A19837) at 1:1000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.




Immunoprecipitation analysis of 300 μ g extracts of K-562 cells using 3 μ g SATB2 antibody (A19837). Western blot was performed from the immunoprecipitate using SATB2 antibody (A19837) at a dilution of 1:1000.


Validation Data



Immunohistochemistry analysis of paraffin-embedded Human appendix using SATB2 Rabbit mAb (A19837) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using SATB2 Rabbit mAb (A19837) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon using SATB2 Rabbit mAb (A19837) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.

