

DNAJC17 Rabbit pAb

Catalog NO.: A20168

Basic Information

Observed MW

35kDa

Calculated MW

35kDa

Category

Polyclonal Antibodies

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Background

Predicted to enable RNA binding activity. Predicted to act upstream of or within negative regulation of transcription by RNA polymerase II and toxin transport. Predicted to be located in cytoplasm and nucleus.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

55192

Swiss Prot

Q9NVM6

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

DNAJC17


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.		


Validation Data




Western blot analysis of various lysates using DNAJC17 Rabbit pAb (A20168) at 1:1000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 180s.



Western blot analysis of various lysates using DNAJC17 Rabbit pAb (A20168) at 1:1000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Enhanced Kit (RM00021).
 Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded Human esophageal cancer using DNAJC17 Rabbit pAb (A20168) at dilution of 1:20 (40x lens). High pressure



Immunohistochemistry analysis of paraffin-embedded Mouse spinal cord using DNAJC17 Rabbit pAb (A20168) at dilution of 1:20 (40x lens). High pressure antigen

antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.