

SATB2 Rabbit mAb

Catalog NO.: A20220 **Recombinant**

Basic Information

Observed MW

100 kDa

Calculated MW

83 kDa

Category

Monoclonal Antibodies

Applications

WB, IP, IF-P, IHC-P, mIHC, ELISA

Cross-Reactivity

Human, Mouse

Clone number

ARC50977

Background

This gene encodes a DNA binding protein that specifically binds nuclear matrix attachment regions. The encoded protein is involved in transcription regulation and chromatin remodeling. Defects in this gene are associated with isolated cleft palate and cognitive disability. Alternate splicing results in multiple transcript variants that encode the same protein.

Recommended Dilutions

WB 1:2000 - 1:10000

IP

0.5 µg - 4 µg antibody for 200 µg - 400 µg extracts of whole cells

IF-P 1:200 - 1:800

IHC-P 1:8000 - 1:40000

mIHC 1:10000 - 1:40000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Contact

Immunogen Information

Gene ID

23314

Swiss Prot

Q9UPW6

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms


GLSS; DEL2Q32Q33; C2DELq32q33; SATB2

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		

Validation Data



Western blot analysis of various lysates using SATB2 Rabbit mAb (A20220) at1:10000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 90s.






Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma(positive) and lung adenocarcinoma(negative) tissue using SATB2 Rabbit mAb (A20220) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human appendix tissue using SATB2 Rabbit mAb (A20220) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using SATB2 Rabbit mAb (A20220) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using SATB2 Rabbit mAb (A20220) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using SATB2 Rabbit mAb (A20220) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

The multiplex IHC analysis on paraffin-embedded Mouse brain tissue using the following specific primary antibodies and tyramide signal amplification (TSA) reagents (RK05903) : SATB2 Rabbit mAb (A20220, 1:20000) with TSA-TYR-520 (Green), and VGLUT1 Rabbit mAb (A26751, 1:500) with TSA-TYR-570 (Red). DAPI (Blue) was used for nuclear staining. Prior to multiplex IHC staining, high-pressure antigen retrieval was performed using 0.01M citrate buffer at pH 6.0. The analysis was completed using a 40x objective lens.

Validation Data



Confocal imaging of paraffin-embedded Mouse brain tissue using SATB2 Rabbit mAb (A20220, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

