

# ASB17 Rabbit pAb

Catalog NO.: A20238

## Basic Information

### Observed MW

34kDa

### Calculated MW

34kDa

### Category

Polyclonal Antibodies

### Applications

WB, ELISA

### Cross-Reactivity

Human, Rat

### Conjugation

Unconjugated

## Background

Predicted to be involved in intracellular signal transduction. Orthologous to human ASB17 (ankyrin repeat and SOCS box containing 17).

## Recommended Dilutions

**WB** 1:500 - 1:1000

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

66772

### Swiss Prot

Q8VHP9

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

4933436O18Rik; ASB17

## Contact

## Product Information

|   |  |                      |
|---|--|----------------------|
| ☒ |  | order@abclonal.com   |
| ☒ |  | service@abclonal.com |
| ☒ |  | www.abclonal.com     |

## Source

Rabbit

## Isotype

IgG

## Purification


Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of lysates from LNCaP cells, using ASB17 Rabbit pAb (A20238) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Western blot analysis of lysates from Rat testis, using ASB17 Rabbit pAb (A20238) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.

