

ECD Rabbit pAb

Catalog NO.: A22237

Basic Information

Observed MW

76kDa

Calculated MW

73kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Human,Mouse,Rat

Conjugation

Unconjugated

Background

Enables histone acetyltransferase binding activity. Involved in positive regulation of transcription by RNA polymerase II. Located in cytosol and nucleoplasm.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

11319

Swiss Prot

O95905

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

GCR2; SGT1; HSGT1; ECD

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from HeLa cells, using ECD Rabbit pAb (A22237) at 1:800 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Western blot analysis of various lysates, using ECD Rabbit pAb (A22237) at 1:800 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.

