

Ly-6G Rabbit mAb

Catalog NO.: A22270 **Recombinant** **9 Publications**

Basic Information

Observed MW

20-40 kDa

Calculated MW

14 kDa

Category

Monoclonal Antibodies

Applications

WB, IF-F, IF-P, IHC-P, ELISA

Cross-Reactivity

Mouse

Conjugation

Unconjugated

Clone number

ARC55430

Background

Located in external side of plasma membrane.

Recommended Dilutions

WB	1:5500 - 1:22000
IF-F	1:200 - 1:800
IF-P	1:400 - 1:1600
IHC-P	1:2000 - 1:8000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Contact

Immunogen Information

Gene ID	Swiss Prot
546644	P35461

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms


Gr1; Gr-1; Ly-6G

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.09% Sodium azide,0.05% BSA,50% glycerol,pH7.3.		

Validation Data



Western blot analysis of various lysates using Ly-6G Rabbit mAb (A22270) at 1:11000 dilution incubated overnight at 4°C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Negative control (NC): NIH/3T3


Exposure time: 45s.





Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using Ly-6G Rabbit mAb (A22270) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of paraffin-embedded Mouse spleen tissue using Ly-6G Rabbit mAb (A22270, dilution 1:800) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of paraffin-embedded Mouse lung tissue using Ly-6G Rabbit mAb (A22270, dilution 1:800) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of frozen sections of Mouse bone marrow tissue using Ly-6G Rabbit mAb (A22270, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Confocal imaging of frozen sections of Mouse spleen tissue using Ly-6G Rabbit mAb (A22270, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Antibody | Product | Research | For Research Use Only

For research use only. Not for therapeutic or diagnostic purposes.

Please visit <https://abclonal.com> for a complete list of recommended products.