

# Uhrf2 Rabbit pAb

Catalog NO.: A2344 **1 Publications**

## Basic Information

### Observed MW

95kDa

### Calculated MW

90kDa

### Category

Polyclonal Antibodies

### Applications

WB, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### Conjugation

Unconjugated

## Background

Enables histone binding activity. Predicted to be involved in several processes, including maintenance of DNA methylation; protein autoubiquitination; and regulation of cell cycle. Located in heterochromatin and nucleus. Is expressed in several structures, including central nervous system; ear; early conceptus; female reproductive system; and urinary system. Orthologous to human UHRF2 (ubiquitin like with PHD and ring finger domains 2).

## Recommended Dilutions

**WB** 1:500 - 1:1000

### ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

109113

### Swiss Prot

Q7TMI3

### Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms

Nirf; 2310065A22Rik; D130071B19Rik; Uhrf2

## Contact

## Product Information

|   |  |                      |
|---|--|----------------------|
| ☒ |  | order@abclonal.com   |
| ☒ |  | service@abclonal.com |
| ☒ |  | www.abclonal.com     |

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of lysates from Mouse thymus, using Uhrf2 Rabbit pAb (A2344) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.

