

S-Tag Rabbit pAb

Catalog NO.: A24465

Basic Information

Observed MW

30kDa

Calculated MW

Category

Tag Antibodies

Applications

WB, ELISA

Cross-Reactivity

Species independent

Conjugation

Unconjugated

Background

S tag is an epitope tag composed of a 15 residue peptide, KETAAAKFERQHMS, derived from the pancreatic ribonuclease A.

Recommended Dilutions

WB 1:1000 - 1:5000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID **Swiss Prot**

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

S-Tag; S

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from wild type (WT) and 293T transfected with S-Tag using S-Tag Rabbit pAb (A24465) at 1:2000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.

