

# SLC1A7 Rabbit pAb

Catalog NO.: A24854

## Basic Information

### Observed MW

70kDa

### Calculated MW

17kDa/60kDa

### Category

Polyclonal Antibodies

### Applications

WB, ELISA

### Cross-Reactivity

Mouse

### Conjugation

Unconjugated

## Background

Predicted to enable anion transmembrane transporter activity. Involved in neurotransmitter reuptake. Predicted to be located in plasma membrane. Predicted to be active in glutamatergic synapse. Predicted to be integral component of postsynaptic membrane and integral component of presynaptic membrane.

## Recommended Dilutions

**WB** 1:500 - 1:1000

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

6512

### Swiss Prot

O00341

### Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms

SLC1A7; AAAT; EAAT5; solute carrier family 1 member 7

## Contact

## Product Information

|   |  |                      |
|---|--|----------------------|
| ☒ |  | order@abclonal.com   |
| ☒ |  | service@abclonal.com |
| ☒ |  | www.abclonal.com     |

## Source

Rabbit

## Isotype

IgG

## Purification


Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of lysates from Mouse testis using SLC1A7 Rabbit pAb (A24854) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.

