

ABflo® 594 Rabbit anti-Human CD27 mAb

Catalog NO.: A25110

Basic Information

Observed MW

Refer to figures

Calculated MW

29kDa

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Human

Conjugation

ABflo® 594. Ex:588nm. Em:604nm.

Clone number

ARC54542

Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.

Recommended Dilutions

FC 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID

939

Swiss Prot

P26842

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

T14; S152; Tp55; TNFRSF7; S152. LPFS2


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		

Validation Data



Flow cytometry:1X106 Human PBMC were surface-stained with ABflo® 488 Rabbit anti-Human/Monkey CD19 mAb (A23008,5 µl/Test) and ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,left) or ABflo® 594 Rabbit anti-Human CD27 mAb(A25110,5 µl/Test,right). Cells in the lymphocyte gate were used for analysis.

