

FITC Rabbit Anti-Goat IgG (Fc) fragment mAb

Catalog NO.: A25513

Basic Information

Observed MW

Calculated MW

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Goat

Conjugation

FITC. Ex:491nm. Em:516nm.

Clone number

ARC56943

Background

Recommended Dilutions

FC 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID Swiss Prot

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		

Validation Data



Flow cytometry: 1X10^6 Jurkat cells (negative control,left) and A-431 cells (right) were surface-stained with Goat anti-Human TROP-2 mAb (IgG, κ,orange line) or FITC Rabbit IgG isotype control (5 μl/Test,blue line), followed by FITC Rabbit Anti-Goat IgG (Fc) fragment mAb (A25513,5 μl/Test) staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10^6 A-431 cells were surface-stained with FITC Rabbit IgG isotype control (5 μl/Test,left) or Goat anti-Human TROP-2 mAb (IgG, κ,right). FITC Rabbit Anti-Goat IgG (Fc) fragment mAb (A25513,5 μl/Test) was used as a secondary antibody.

