

PE Rabbit Anti-Goat IgG (Fc) fragment mAb

Catalog NO.: A25517

Basic Information

Observed MW**Calculated MW**

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Goat

Conjugation

PE. Ex:565nm. Em:574nm.

Clone number

ARC56943

Background

Recommended Dilutions

FC 5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage Store at 2-8°C. Avoid freeze. Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		

Validation Data



Flow cytometry: 1X10⁶ Jurkat cells (negative control,left) and A-431 cells (right) were surface-stained with Goat anti-Human TROP-2 mAb (IgG, κ,orange line) or PE Rabbit IgG isotype control (A24172,5 μl/Test,blue line), followed by PE Rabbit Anti-Goat IgG (Fc) fragment mAb (A25517,5 μl/Test) staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10⁶ A-431 cells were surface-stained with PE Rabbit IgG isotype control (A24172,5 μl/Test,left) or Goat anti-Human TROP-2 mAb (IgG, κ,right). PE Rabbit Anti-Goat IgG (Fc) fragment mAb (A25517,5 μl/Test) was used as a secondary antibody.

