

PerCP/Cyanine5.5 Rabbit IgG isotype control

Catalog NO.: A25620

Basic Information

Observed MW

Calculated MW

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Conjugation

PerCP-Cy5.5. Ex:482nm. Em:695nm.

Clone number

ARC5105-10

Background

The isotype of a primary antibody and the application it is being used in can result in background staining. Primary antibody background noise can be caused by binding to Fc receptors on target cells; by non-specific interactions with cellular proteins, carbohydrates, and lipids; or by cell autofluorescence. Isotype control antibodies can act as negative controls to help differentiate non-specific background signal from specific antibody signal because they have no relevant specificity to a target antigen. An isotype control antibody should have the same immunoglobulin type and be used at the same concentration as the test antibody.

Recommended Dilutions

FC 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Flow cytometry: 1X10⁶ 293T cells (negative control, left) and Jurkat cells (right) were surface-stained with PerCP/Cyanine5.5 Rabbit anti-Human CD45 mAb (A25622, 5 µl/Test, orange line) or PerCP/Cyanine5.5 Rabbit IgG isotype control (A25620, 5 µl/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

