

Arginase-1 Rabbit mAb

Catalog NO.: A25808 **Recombinant** **18 Publications**

Basic Information

Observed MW

40kDa

Calculated MW

35kDa

Category

Monoclonal Antibodies

Applications

WB, IF-P, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Clone number

ARC65709

Background

Enables arginase activity. Involved in defense response to protozoan; negative regulation of T-helper 2 cell cytokine production; and negative regulation of activated T cell proliferation. Predicted to be located in several cellular components, including extracellular space; mitochondrial outer membrane; and neuronal cell body. Predicted to be active in cytosol. Is expressed in several structures, including alimentary system; central nervous system; genitourinary system; integumental system; and sensory organ. Used to study hyperargininemia. Human ortholog(s) of this gene implicated in asthma; hepatocellular carcinoma; and hyperargininemia. Orthologous to human ARG1 (arginase 1).

Recommended Dilutions

WB 1:10000 - 1:40000

IF-P 1:200 - 1:800

IHC-P 1:2000 - 1:10000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID	Swiss Prot
11846	Q61176

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

AI; PGIF; Arg-1; Arginase-1


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.09% sodium azide,0.05% BSA,50% glycerol,pH7.3.		

Validation Data



Western blot analysis of various lysates using Arginase-1 Rabbit mAb (A25808)at 1:10000 dilution incubated at room temperature for 1.5 hours.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Negative control (NC): Mouse small intestine


Exposure time: 20s.





Immunohistochemistry analysis of paraffin-embedded Human liver tissue using Arginase-1 Rabbit mAb (A25808) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using Arginase-1 Rabbit mAb (A25808) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat liver tissue using Arginase-1 Rabbit mAb (A25808) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of paraffin-embedded Human liver tissue using Arginase-1 Rabbit mAb (A25808, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Antibody Product Q&A with Enzo Life Sciences

Confocal imaging of paraffin-embedded Mouse liver tissue using Arginase-1 Rabbit mAb (A25808, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit <https://abclonal.com> for a complete list of recommended products.