

# ICAM5 Rabbit mAb

Catalog NO.: A26618 **Recombinant** **1 Publications**

## Basic Information

### Observed MW

150kDa

### Calculated MW

97kDa

### Category

Monoclonal Antibodies

### Applications

WB, IF-P, IHC-P, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### Conjugation

Unconjugated

### Clone number

ARC69846

## Background

Predicted to enable integrin binding activity. Acts upstream of or within phagocytosis. Located in plasma membrane. Is expressed in brain. Orthologous to human ICAM5 (intercellular adhesion molecule 5).

## Recommended Dilutions

**WB** 1:50000 - 1:300000

**IF-P** 1:400 - 1:1600

**IHC-P** 1:1000 - 1:4000

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

15898

### Swiss Prot

Q60625

### Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms

TLN; CD50; Tlcn; Icam3; ICAM-3; ICAM-5


## Contact

## Product Information

|                                     |  |                      |
|-------------------------------------|--|----------------------|
| <input checked="" type="checkbox"/> |  | order@abclonal.com   |
| <input checked="" type="checkbox"/> |  | service@abclonal.com |
| <input checked="" type="checkbox"/> |  | www.abclonal.com     |

|                                                                   |                |                       |
|-------------------------------------------------------------------|----------------|-----------------------|
| <b>Source</b>                                                     | <b>Isotype</b> | <b>Purification</b>   |
| Rabbit                                                            | IgG            | Affinity purification |
| <b>Storage</b>                                                    |                |                       |
| Store at -20°C. Avoid freeze / thaw cycles.                       |                |                       |
| Buffer: PBS with 0.09% Sodium azide,0.05% BSA,50% glycerol,pH7.3. |                |                       |

## Validation Data



Western blot analysis of various lysates using ICAM5 Rabbit mAb (A26618) at 1:50000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.




Immunohistochemistry analysis of paraffin-embedded Human brain tissue using ICAM5 Rabbit mAb (A26618) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using ICAM5 Rabbit mAb (A26618) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using ICAM5 Rabbit mAb (A26618) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.



Confocal imaging of paraffin-embedded Human brain tissue using ICAM5 Rabbit mAb (A26618, dilution 1:400) followed by a further incubation with Cy3 Goat Anti-



Confocal imaging of paraffin-embedded Rat brain tissue using ICAM5 Rabbit mAb (A26618, dilution 1:400) followed by a further incubation with Cy3 Goat Anti-



Confocal imaging of paraffin-embedded Mouse brain tissue using ICAM5 Rabbit mAb (A26618, dilution 1:400) followed by a further incubation with Cy3 Goat Anti-

Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IF staining. Objective: 40x.

Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IF staining. Objective: 40x.

Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IF staining. Objective: 40x.