

PE Rabbit anti-Mouse NCAM mAb

Catalog NO.: A29003

Basic Information

Observed MW

Calculated MW

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Mouse

Conjugation

PE. Ex:565nm. Em:574nm.

Clone number

ARC60214

Background

Predicted to enable LRR domain binding activity and phosphatase binding activity. Involved in commissural neuron axon guidance and regulation of semaphorin-plexin signaling pathway. Acts upstream of or within several processes, including homotypic cell-cell adhesion; positive regulation of calcium-mediated signaling; and regulation of exocyst assembly. Located in several cellular components, including external side of plasma membrane; growth cone; and neuronal cell body. Is expressed in several structures, including embryo mesenchyme; nervous system; sensory organ; skin; and urinary system. Human ortholog(s) of this gene implicated in bipolar disorder; middle cerebral artery infarction; and pancreatic cancer. Orthologous to human NCAM1 (neural cell adhesion molecule 1).

Recommended Dilutions

FC ≤0.5 µg per million cells in 100 µl volume

Immunogen Information

Gene ID

17967

Swiss Prot

P13595

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

CD56; Ncam; E-NCAM; NCAM-1


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.		

Validation Data



Flow cytometry: 1X10⁶ BALB/c mouse bone marrow cells were surface-stained with PE Rabbit IgG isotype control (A24172,5 µl/Test,left) or PE Rabbit anti-Mouse NCAM mAb (A29003,0.5 µg,right). Total viable cells were used for analysis.

