

ABflo® 450 Rabbit anti-Human/Monkey CD16 mAb

Catalog NO.: A29409

Basic Information

Observed MW

Calculated MW

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Human

Conjugation

ABflo® 450. Ex:406nm. Em:445nm.

Clone number

ARC59985

Background

This gene encodes a receptor for the Fc portion of immunoglobulin G, and it is involved in the removal of antigen-antibody complexes from the circulation, as well as other responses, including antibody dependent cellular mediated cytotoxicity and antibody dependent enhancement of virus infections. This gene (FCGR3A) is highly similar to another nearby gene (FCGR3B) located on chromosome 1. The receptor encoded by this gene is expressed on natural killer (NK) cells as an integral membrane glycoprotein anchored through a transmembrane peptide, whereas FCGR3B is expressed on polymorphonuclear neutrophils (PMN) where the receptor is anchored through a phosphatidylinositol (PI) linkage. Mutations in this gene are associated with immunodeficiency 20, and have been linked to susceptibility to recurrent viral infections, susceptibility to systemic lupus erythematosus, and alloimmune neonatal neutropenia. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

FC 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID

Hu 2214/2215/Cyno 102140945

Swiss Prot

P08637/O75015

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

CD16; FCG3; CD16A; FCGR3; IGFR3; IMD20; FCR-10; FCRIII; CD16-II; FCGRIII; FCRIIIA; FcGRIIIA


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.		

Validation Data



Flow cytometry: 1X10⁶ Human PBMC were surface-stained with ABflo® 450 Rabbit IgG isotype control (A27451,5 µl/Test,left) or ABflo® 450 Rabbit anti-Human/Monkey CD16 mAb (A29409,5 µl/Test,right). Cells in the lymphocyte gate were used for analysis.



Flow cytometry: 1X10⁶ Cynomolgus PBMC were surface-stained with ABflo® 450 Rabbit IgG isotype control (A27451,5 µl/Test,left) or ABflo® 450 Rabbit anti-Human/Monkey CD16 mAb (A29409,5 µl/Test,right). Cells in the lymphocyte gate were used for analysis.

