

# EIF3M Rabbit pAb

Catalog NO.: A4426

## Basic Information

### Observed MW

40kDa/36kDa

### Calculated MW

42kDa

### Category

Polyclonal Antibodies

### Applications

WB, IHC-P, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### Conjugation

Unconjugated

## Background

This gene encodes a protein that is part of the eukaryotic translation initiation factor 3 complex (eIF-3) required for protein synthesis. Elevated levels of the encoded protein are present in cancer cell lines. Inactivation of the encoded protein has been shown to interfere with translation of herpes virus mRNAs by preventing the association of mRNAs with the ribosomes. A pseudogene of this gene is located on the X chromosome.

## Recommended Dilutions

**WB** 1:500 - 1:2000

**IHC-P** 1:50 - 1:200

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

10480

### Swiss Prot

Q7L2H7

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

B5; GA17; PCID1; TANGO7; hfl-B5; EIF3M


## Contact

## Product Information

|                                     |  |                      |
|-------------------------------------|--|----------------------|
| <input checked="" type="checkbox"/> |  | order@abclonal.com   |
| <input checked="" type="checkbox"/> |  | service@abclonal.com |
| <input checked="" type="checkbox"/> |  | www.abclonal.com     |

|                                                                                                                                       |                |                       |
|---------------------------------------------------------------------------------------------------------------------------------------|----------------|-----------------------|
| <b>Source</b>                                                                                                                         | <b>Isotype</b> | <b>Purification</b>   |
| Rabbit                                                                                                                                | IgG            | Affinity purification |
| <b>Storage</b>                                                                                                                        |                |                       |
| Store at -20°C. Avoid freeze / thaw cycles.                                                                                           |                |                       |
| Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3. |                |                       |

Validation Data



Western blot analysis of various lysates using EIF3M Rabbit pAb (A4426) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Western blot analysis of various lysates using EIF3M Rabbit pAb (A4426) at 1:1000 dilution incubated overnight at 4°C.



Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020)

Exposure time: 5 s.



Immunohistochemistry analysis of paraffin-embedded Human liver damage using EIF3M Rabbit pAb (A4426) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Rat brain using EIF3M Rabbit pAb (A4426) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.