INTS4 Rabbit pAb

Catalog NO.: A6339

Basic Information

Observed MW
108kDa

Calculated MW
108kDa

Category
Polyclonal Antibodies

Applications
WB, IP, ELISA

Cross-Reactivity
Human, Mouse, Rat

Conjugation
Unconjugated

Background

INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]).

Recommended Dilutions

WB 1:500 - 1:1000

IP
0.5μg-4μg antibody for 200μg-400μg extracts of whole cells

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
92105

Swiss Prot
Q96HW7

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
INT4; MST093; INTS4

Contact

Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Western blot analysis of various lysates using INTS4 Rabbit pAb (A6339) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.



Western blot analysis of lysates from Rat liver, using INTS4 Rabbit pAb (A6339) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Immunoprecipitation analysis of 300 µg extracts of Jurkat cells using 3 µg INTS4 antibody (A6339). Western blot was performed from the immunoprecipitate using INTS4 antibody (A6339) at a dilution of 1:500.