

FKBP3 Rabbit pAb

Catalog NO.: A6907

Basic Information

Observed MW

30kDa

Calculated MW

25kDa

Category

Polyclonal Antibodies

Applications

WB, IP, ELISA

Cross-Reactivity

Human

Conjugation

Unconjugated

Background

The protein encoded by this gene is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. This encoded protein is a cis-trans prolyl isomerase that binds the immunosuppressants FK506 and rapamycin, as well as histone deacetylases, the transcription factor YY1, casein kinase II, and nucleolin. It has a higher affinity for rapamycin than for FK506 and thus may be an important target molecule for immunosuppression by rapamycin.

Recommended Dilutions

WB 1:500 - 1:2000

IP

0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

2287

Swiss Prot

Q00688

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

FKBP-3; FKBP25; PPIase; FKBP-25; FKBP3

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates using FKBP3 Rabbit pAb (A6907) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 30s.



Immunoprecipitation analysis of 200 µg extracts of MCF-7 cells, using 3 µg FKBP3 antibody (A6907). Western blot was performed from the immunoprecipitate using FKBP3 antibody (A6907) at a dilution of 1:1000.

