

TRMT1 Rabbit pAb

Catalog NO.: A7116

Basic Information

Observed MW

90kDa

Calculated MW

72kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse

Conjugation

Unconjugated

Background

This gene encodes a tRNA-modifying enzyme that acts as a dimethyltransferase, modifying a single guanine residue at position 26 of the tRNA. The encoded enzyme has both mono- and dimethylase activity when exogenously expressed, and uses S-adenosyl methionine as a methyl donor. The C-terminal region of the encoded protein has both a zinc finger motif, and an arginine/proline-rich region. Mutations in this gene have been implicated in autosomal recessive intellectual disorder (ARID). Alternative splicing results in multiple transcript variants encoding different isoforms. There is a pseudogene of this gene on the X chromosome.

Recommended Dilutions

WB

1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

55621

Swiss Prot

Q9NXH9

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

TRM1; MRT68; TRMT1

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates using TRMT1 Rabbit pAb (A7116) at 1:1000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Enhanced Kit (RM00021).
 Exposure time: 90s.

