

ERCC4 Rabbit pAb

Catalog NO.: A8119 **2 Publications**

Basic Information

Observed MW

120kDa

Calculated MW

104kDa

Category

Polyclonal Antibodies

Applications

WB, IP, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Background

The protein encoded by this gene forms a complex with ERCC1 and is involved in the 5' incision made during nucleotide excision repair. This complex is a structure specific DNA repair endonuclease that interacts with EME1. Defects in this gene are a cause of xeroderma pigmentosum complementation group F (XP-F), or xeroderma pigmentosum VI (XP6).

Recommended Dilutions

WB 1:500 - 1:2000

IP

0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

2072

Swiss Prot

Q92889

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

XPF; RAD1; FANCO; XFEPS; ERCC11; ERCC4

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates using ERCC4 Rabbit pAb (A8119) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Immunoprecipitation analysis of 150 µg extracts of 293T cells using 3 µg ERCC4 antibody (A8119). Western blot was performed from the immunoprecipitate using ERCC4 antibody (A8119) at a dilution of 1:1000.

