

Pan Phospho-Threonine Rabbit mAb

Catalog NO.: AP1422 **Recombinant** **1 Publications**

Basic Information

Observed MW

10-250kDa/

Calculated MW

Category

Phosphorylated Antibodies

Applications

WB, IP, ELISA

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

Conjugation

Unconjugated

Clone number

ARC61310

Background

Recommended Dilutions

WB 1:500 - 1:1000

IP

0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID **Swiss Prot**

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from A-431, using Pan Phospho-Threonine Rabbit mAb (AP1422) at 1:1000 dilution. A 431 treated with Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 60s.



Western blot analysis of lysates from NIH 3T3 or NIH 3T3 treat with Calyculin A, using Pan Phospho-Threonine Rabbit mAb (AP1422) at 1:1000 dilution. NIH/3T3 cells were treated with Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 60s.



Immunoprecipitation of Pan Phospho-Threonine from 300 µg extracts of A-431 cells treated with calyculin A (200nM,30min) was performed using 2 µg of Pan Phospho-Threonine Rabbit mAb (AP1422). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1x reducing Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using Pan Phospho-Threonine Rabbit mAb (AP1422) at a dilution of 1:1000.