

HRP-conjugated Mouse anti-Rabbit IgG (H+L)

Catalog No.: AS079**4 Publications**

Basic Information

Observed MW

36kDa

Calculated MW**Category**

Secondary Antibodies

Applications

WB, ELISA

Cross Reactivity

Rabbit

Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies. Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

Recommended Dilutions

WB

1:5000 - 1:20000

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID**Swiss-Prot****Immunogen**

This information is considered to be commercially sensitive.

Synonyms

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Mouse

Isotype

IgG



Purity

Affinity purification

Storage

Store at -20°C

Validation Data



Dose response curve of HRP-conjugated Mouse anti-Rabbit IgG (H+L) measured by ELISA. 1 μ g/mL of various immunoglobulins were coated to 384-well plate. The coated plate was blocked and subsequently incubated with 25 μ L of HRP conjugated in a 3 fold serial dilution from 1 μ g/mL to 5.2×10^{-3} μ g/mL, incubation was performed at room temperature for 1 hour. The ELISA result demonstrated that HRP-conjugated Mouse anti-Rabbit IgG (H+L) has highly specific recognition of Rabbit IgG while no or minimal cross reactivity to Human IgG, Mouse IgG, Rat IgG, Bovine IgG, Goat IgG.