

HRP conjugated Rabbit anti-Mouse IgM (Fc) mAb

Catalog NO.: AS094 **1 Publications**

Basic Information

Observed MW

55kDa

Calculated MW

Category

Secondary Antibodies

Applications

WB, ELISA

Cross-Reactivity

Mouse

Conjugation

HRP

Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies . Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

Recommended Dilutions

WB 1:10000 - 1:50000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID **Swiss Prot**

Immunogen

This information is considered to be commercially sensitive.

Synonyms


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit		Affinity purification
Storage		
Store at -20°C. Avoid freeze.		
Buffer: PBS containing 50% glycerol and 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		

Validation Data



Western blot analysis of lysates from Mouse IgM using HRP conjugated Rabbit anti-Mouse IgM (Fc) mAb (AS094) at 1:50000 dilution.

Lysates/proteins: 50ng per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45s.

