

HRP-conjugated Rabbit anti-Goat IgG Fab mAb

Catalog NO.: AS097 **1 Publications**

Basic Information

Observed MW

Calculated MW

Category

Secondary Antibodies

Applications

WB, ELISA

Cross-Reactivity

Goat

Conjugation

HRP

Clone number

ARC57078-HRP

Background

Recommended Dilutions

WB 1:500-1:64000

ELISA 1:5000-1:10000

Immunogen Information

Gene ID **Swiss Prot**

Immunogen

This information is considered to be commercially sensitive.

Synonyms


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		

Validation Data




Western blot analysis of recombinant goat IgG Fab protein using HRP-conjugated Rabbit anti-Goat IgG Fab mAb(AS097) at 1:500-1:64000 dilution. Blot was incubated with the HRP conjugated antibody in the Blocking buffer (3% non-fat dry milk in TBST) at room temperature for 1 hour before detection using Enhanced chemiluminescence.


Lysates/proteins: Lane 1: molecular weight marker; Lanes 2-9: 100ng goat IgG Fab recombinant Protein per lane.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s



Dose response curve of HRP conjugated Rabbit Anti-Goat IgG Fab mAb measured by ELISA. 25 ng of Goat IgG Fab recombinant protein was coated in 384-well plate, blank wells without protein were used as negative control (NC). 25 μ L of HRP conjugated Rabbit Anti-Goat IgG Fab mAb in a 2 fold serial dilution from 1 μ g/mL to 7.8 pg/mL, incubation was performed at room temperature for 1 hour. EC50 value (4.27×10^{-2} μ g/mL) were calculated using 4 parameter logistic regression.



Cross reactivity to various IgGs of HRP conjugated Rabbit Anti-Goat IgG Fab mAb was examined ELISA. 1 μ g/mL of various immunoglobulins were coated to 384-well plate. Then the plate was blocked and incubated with 25 μ L of HRP conjugated Rabbit Anti-Goat IgG Fab mAb , 0.5 μ g/mL and 0.125 μ g/mL) at room temperature for 1 hour. The ELISA result demonstrated that Rabbit anti-goat Fab antibody has highly specific recognition of goat IgG (H+L) and Fab fragments, while no or minimal cross reactivity to goat IgG Fc, Human IgG, Swine IgG, Horse IgG, Chicken IgY, and Rat IgG.

