

ABflo® 647 Rabbit anti-Rat IgG2b (Fc) mAb

Catalog NO.: AS165

Basic Information

Observed MW

Calculated MW

36kDa

Category

Secondary Antibodies

Applications

FC

Cross-Reactivity

Rat

Conjugation

ABflo® 647. Ex:648nm. Em:664nm.

Clone number

ARC72127-ABflo647

Background

Enables antigen binding activity. Acts upstream of or within phagocytosis; positive regulation of immune response; and positive regulation of phagocytosis. Located in extracellular space. Part of immunoglobulin complex, circulating. Orthologous to several human genes including IGHG1 (immunoglobulin heavy constant gamma 1 (G1m marker)); IGHG2 (immunoglobulin heavy constant gamma 2 (G2m marker)); and IGHG3 (immunoglobulin heavy constant gamma 3 (G3m marker)).

Recommended Dilutions

FC 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID

Swiss Prot

P20761

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

IgG2b; Igh-3; gamma2b


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.		

Validation Data



Flow cytometry: 1X10⁶ Neuro-2a cells (negative control,left) and C57BL/6 mouse splenocytes (right) were surface-stained with Purified Rat anti-mouse I-A/I-E Antibody (2ug/ml,orange line) or Rat IgG2b isotype control (2 μg/mL,blue line), followed by ABflo® 647 Rabbit anti-Rat IgG2b (Fc) mAb (AS165,5 μl/Test) staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10⁶ C57BL/6 mouse splenocytes were surface-stained with Rat IgG2b isotype control (2 μg/mL,left) or Purified Rat anti-mouse I-A/I-E Antibody (2ug/ml,right), followed by ABflo® 647 Rabbit anti-Rat IgG2b (Fc) mAb (AS165,5 μl/Test) staining.

