

Recombinant Human IL-22RA2/IL22BP Protein

Catalog No.: RP00192

Sequence Information

Species **Gene ID**
Human 116379

Swiss Prot **Tags**
Q969J5-2 C-His

Synonyms

IL22RA2, CRF2-10, CRF2-S1, CRF2X,
IL-22BP, IL-22R-alpha-2, IL-22RA2,
ZCYTOR16

Product Information

Source **Purification**
HEK293 cells ≥ 90 % as
 determined by
 SDS-PAGE.

Calculated MW **Observed MW**
25.33 kDa 40-50 kDa

Endotoxin
< 0.1 EU/μg of the protein by LAL
method.

Formulation

Lyophilized from a 0.22 μm filtered
solution of PBS, pH 7.4. Contact us
for customized product form or
formulation.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of
0.1-0.5 mg/mL in sterile distilled
water. Avoid vortex or vigorously
pipetting the protein. For long
term storage, it is recommended
to add a carrier protein or stabilizer
(e.g. 0.1% BSA, 5% HSA, 10% FBS
or 5% Trehalose), and aliquot the
reconstituted protein solution to
minimize free-thaw cycles.

Background

Interleukin 22 binding protein (IL-22BP), also known as CRF2-10, CRF2-X, and IL-22RA2, is a 35-45 kDa secreted glycoprotein in the type II cytokine receptor family (CRF). IL-22BP specifically binds to and inhibits interleukin 22 activity by blocking the interaction of interleukin 22 with its cell surface receptor. IL-22BP is produced by dendritic cells (DC), epithelial cells, activated B cells, and activated monocytes. It is constitutively expressed by DC but is down-regulated during local inflammation and in response to tissue damage. IL-22BP is critical for limiting IL-22 induced epithelial cell proliferation during wound healing, and its deficiency can enable uncontrolled proliferation and enhance tumor development.

Basic Information

Description

Recombinant Human IL-22RA2/IL22BP Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Thr22-Pro231) of human IL-22BP/IL-22RA2 (Accession #NP_851826.1) fused with a 6×His tag at the C-terminus.

Activity

Measured by its binding ability in a functional ELISA. Immobilized recombinant Human IL22 at 2 μg/mL (100 μL/well) can bind recombinant Human IL22BP, the EC₅₀ of Human IL22BP is 10.78 ng/mL.

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Recombinant Human IL-22RA2/IL22BP Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized recombinant Human IL22 at 2µg/mL (100µL/well) can bind recombinant Human IL22BP, the EC₅₀ of Human IL22BP is 10.78 ng/mL.