

Recombinant Mouse CX3CL1/Fractalkine Protein

Catalog No.: RP01393

Sequence Information

Species Gene ID

Mouse 20312

Swiss Prot Tags

O35188 C-His

Synonyms

CX3CL1, Fractalkine, Cx3cl1, Cx3c, Fkn, Scyd1, CX3CL1

Product Information

Source Purification

HEK293 cells ≥ 95 % as determined by SDS-PAGE.

Calculated MW Observed MW

34.06 kDa 50–70 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1–0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Fractalkine or Chemokine (C-X3-C motif) ligand 1 (CX3CL1) is a member of the CX3C chemokine family. Fractalkine / CX3CL1 is a unique chemokine that functions not only as a chemoattractant but also as an adhesion molecule and is expressed on endothelial cells activated by proinflammatory cytokines, such as interferon-gamma and tumor necrosis factor-alpha. Fractalkine / CX3CL1 is expressed in a membrane-bound form on activated endothelial cells and mediates attachment and firm adhesion of T cells, monocytes and NK cells. Fractalkine / CX3CL1 is associated with dendritic cells (DC) in epidermis and lymphoid organs. The fractalkine receptor, CX3CR1, is expressed on cytotoxic effector lymphocytes, including natural killer (NK) cells and cytotoxic T lymphocytes, which contain high levels of intracellular perforin and granzyme B, and on macrophages. Soluble fractalkine causes migration of NK cells, cytotoxic T lymphocytes, and macrophages, whereas the membrane-bound form captures and enhances the subsequent migration of these cells in response to secondary stimulation with other chemokines.

Basic Information

Description

Recombinant Mouse CX3CL1/Fractalkine Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln25–Arg337) of mouse CX3CL1/Fractalkine (Accession #NP_033168.2) fused with a 6×His tag at the C-terminus.

Activity

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at –20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Recombinant Mouse CX3CL1/Fractalkine Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.