

Recombinant Human TNFSF12/TWEAK Protein

Catalog No.: RP01446

Sequence Information

Species **Gene ID**
Human 8742

Swiss Prot **Tags**
O43508-1 No tag

Synonyms

TNFSF12, APO3L, DR3LG, TNLG4A, TWEAK

Product Information

Source **Purification**
E. coli ≥ 95 % as
determined by
SDS-PAGE.

Calculated MW **Observed MW**
16.94 kDa 15-20 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

TNFSF12 is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. It is a ligand for the FN14/TWEAKR receptor. TNFSF12 has overlapping signaling functions with TNF, but displays a much wider tissue distribution. It can induce apoptosis via multiple pathways of cell death in a cell type-specific manner. It is also found that TNFSF12 promotes proliferation and migration of endothelial cells, and thus acts as a regulator of angiogenesis. TNFSF12 also is a weak inducer of apoptosis in some cell types and mediates NF-κB activation.

Basic Information

Description

Recombinant Human TNFSF12/TWEAK Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Lys97-His249 (N-Met)) of human TWEAK/TNFSF12 (Accession #NP_003800.1) fused with no additional amino acid.

Activity

1. Measured by its binding ability in a functional ELISA. Immobilized Human TNFSF12 at 2 μg/mL (100 μL/well) can bind Human TNFRSF12A with a linear range of 0.1-2.3 ng/mL.
2. Measured in a cell proliferation assay using human umbilical vein endothelial cells (HUVEC). The ED₅₀ for this effect is typically 0.81-3.24 ng/mL.

Shipping

Lyophilized proteins are shipped at ambient temperature. Liquid proteins are shipped at frozen temperature with dry ice. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Immobilized Human TNFSF12 at 2 µg/mL (100 µL/well) can bind Human TNFRSF12A with a linear range of 0.1–2.3 ng/mL.



Recombinant Human TNFSF12/TWEAK Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Human TNFSF12 stimulates cell proliferation of the human umbilical vein endothelial cells (HUVEC).The ED₅₀ for this effect is typically 0.81–3.24 ng/mL.