Recombinant Human PTPRC/CD45RA/CD45 Protein

Catalog No.: RP01529

Sequence Information

Species
Human

Gene ID
5788

Swiss Prot
P08575-8

Tags
C-Rabbit Fc

Synonyms
B220,CD45,CD45R,GP180,L-CA, LCA,LY5,T200,PTPRC, B220, CD45, CD45R, GP180, L-CA, LCA, LY5, T200, receptor-type tyrosine-protein phosphatase C

Product Information

Source
HEK293 cells

Purification
≥ 95 % as determined by SDS-PAGE.

Calculated MW
76.96 kDa

Observed MW
100-150 kDa

Endotoxin
< 0.1 EU/μg of the protein by LAL method.

Formulation
Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution
Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Protein tyrosine phosphatase, receptor type C (CD45), also known as PTPRC is a member of the protein tyrosine phosphatase (PTP) family which is known for its function to serve as signaling molecules and to regulate a variety of cellular processes such as cell proliferation, differentiation, mitotic cycle and oncogenic transformation. CD45 is found expression specifically in hemotopietic cells. CD45 consists of an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains. It serves as an essential regulator of T-cell and B-cell antigen receptor signaling through either direct interaction with components of the antigen receptor complexes or by activating various Src family kinases required for the antigen receptor signaling and it also can suppress JAK kinases.

Basic Information

Description
Recombinant Human PTPRC/CD45RA/CD45 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln26-Lys482) of human PTPRC/CD45/CD45RA (Accession #AAD15273.2) fused with a raFc tag at the C-terminus.

Activity

Shipping
Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage
Store at -20℃


Operational Notes
For your safety and health, please wear a lab coat and disposable gloves for handling.

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.

Validation Data



Recombinant Human PTPRC/CD45RA/CD45 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.