

Recombinant Rat CXCL1/GRO- α Protein

Catalog No.: RP01917

Sequence Information

Species	Gene ID
Rat	81503
Swiss Prot	Tags
P14095	C-Avi-His

Synonyms

Gro1; CINC-1, rRtGRO- α /CXCL1;
Growth-regulated alpha protein;
C-X-C motif chemokine 1; CINC-1

Product Information

Source	Purification
HEK293 cells	$\geq 95\%$ as determined by SDS-PAGE.

Calculated MW	Observed MW
10.56 kDa	10–15 kDa

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1–0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

CXCL1, also known as KC, GRO α , and CINC-1, is an approximately 8 kDa proinflammatory chemokine that plays a key role in neutrophil migration and activation. Mature human CXCL1 shares 64% and 67% aa sequence identity with mouse and rat CXCL1, respectively. It is produced by many cell types in inflammatory sites and during chronic inflammatory diseases. CXCL1 can associate into bioactive dimers and primarily signals through CXCR2/IL-8 RB but can also bind with lower affinity to CXCR2/IL-8 RA. It induces neutrophil migration, extravasation, respiratory burst, and degranulation and also induces T cells to produce proinflammatory IL-17. CXCL1 additionally binds to Syndecan-1 on epithelial cells which acts as a sink for CXCL1 activity until Syndecan-1 cleavage by MMP-7. CXCL1 is up-regulated in spinal cord astrocytes by inflammatory stimuli or tumor cell injection, and it exacerbates pain sensation by potentiating excitatory NMDA neurotransmission. In the circulatory system, CXCL1 interacts with CXCR2 on endothelial cells to promote lymphatic tube formation and angiogenesis. It promotes the hypertrophic differentiation of chondrocytes resulting in cartilage matrix deposition, calcification, and remodeling. It interacts with both CXCR1 and CXCR2 on adipose stromal cells and promotes their recruitment to prostate tumors in obese patients. It also binds CXCR2 on ovarian cancer cells, leading to cleavage of cell surface HB-EGF, transactivation of EGF R, and cell proliferation.

Basic Information

Description

Recombinant Rat CXCL1/GRO- α Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly24–Lys96) of Rat CXCL1/GRO- α (Accession #NP_110472.1) fused with a Avi-His tag at the C-terminus.

Activity

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

		400-999-6126
		cn.market@abclonal.com.cn
		www.abclonal.com.cn

Validation Data



Recombinant Rat CXCL1/GRO- α Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Measured by its binding ability in a functional ELISA. Immobilized recombinant Rat CXCL1 Antibody at 0.5 $\mu\text{g/mL}$ (25 μL /well) can bind recombinant Rat CXCL1 with a linear range of 0.46–3.43 ng/mL.