

Recombinant Human HSP70 Protein

Catalog No.: RP02123

Sequence Information

Species **Gene ID**
Human 3303

Swiss Prot **Tags**
P0DMV8 N-6His

Synonyms

HSPA1A, HSP72, HSPA1, HSX70,
Heat shock 70 kDa protein 1

Product Information

Source **Purification**
HEK293 cells ≥ 90 % as
determined by
SDS-PAGE.

Calculated MW **Observed MW**
71.4 kDa 70-80 kDa

Endotoxin

< 0.01 EU/μg of the protein by LAL
method.

Formulation

Lyophilized from 0.22μm filtered
solution in PBS (pH 7.4). 5 %
trehalose, 0.01% Tween80 are
added as protectants before
lyophilization. Please contact us for
any concerns or special
requirements.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of
0.1-0.5 mg/mL in sterile distilled
water. Avoid vortex or vigorously
pipetting the protein. For long
term storage, it is recommended
to add a carrier protein or stabilizer
(e.g. 0.1% BSA, 5% HSA, 10% FBS
or 5% Trehalose), and aliquot the
reconstituted protein solution to
minimize free-thaw cycles.

Background

HSPA1A is a member of the Hsp70 protein family. The 70 kilodalton heat shock proteins (Hsp70s) are a family of ubiquitously expressed heat shock proteins. HSP are abundant and conserved proteins present in all cells. Upon temperature shock or other stress stimuli, HSP is synthesized intracellularly, which may protect cells from protein denaturation or death. Extracellularly, HSP can serve a cytokine function to initiate both innate and adaptive immunity through activation of APC. HSP serves also a chaperone function and facilitates the presentation of antigen peptide to T cells. Molecular chaperones of the Hsp70 family have diverse functions in cells. They assist the folding of newly synthesized and stress-denatured proteins, as well as the import of proteins into organelles, and the dissociation of aggregated proteins. The well-conserved Hsp70 chaperones are ATP dependent: binding and hydrolysis of ATP regulate their interactions with unfolded polypeptide substrates, and ATPase cycling is necessary for their function. All cellular functions of Hsp70 chaperones use the same mechanism of ATP-driven polypeptide binding and release.

Basic Information

Description

Recombinant Human HSP70 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala2-Asp641) of Human HSP70 (Accession #NP_005336.3) fused with His tag at the N-Terminus.

Activity

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Recombinant Human HSP70 Protein was determined by SDS-PAGE under reducing (R) and non-reducing (NR) conditions.