

Recombinant Human IL-17Rc Protein

Catalog No.: RP02624

Sequence Information

Species	Gene ID
Human	84818
Swiss Prot	Tags
NP_703190.2	C-hFc

Synonyms

IL-17 receptor C; IL-17RC;
IL17Rhom; IL-17RL; ZcytoR14

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by Tris-Bis PAGE; ≥ 95 % as determined by HPLC.

Calculated MW Observed MW

76.2 kDa 90-115 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

IL-17RC (interleukin-17 receptor-like) gene codes for a transmembrane protein, the full length of which inhibits apoptosis in prostate cancer cells. IL-17RC gene transcribes over a dozen different splice variants of mRNA. IL-17RC protein isoforms are differentially expressed in prostatic cells and cancer tissues and may play a negative or positive role in the initiation and progression of prostate cancer.

Basic Information

Description

Recombinant Human IL-17Rc Protein is expressed from Expi293 with hFc tag at the C-terminal.; It contains Leu21-Arg467.

Activity

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Immobilized Human IL-17A, His Tag at 1µg/ml (100µl/Well) on the plate. Dose response curve for Human IL-17Rc, hFc Tag with the EC₅₀ of 43.5ng/ml determined by ELISA.



Recombinant Human IL-17Rc Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Human IL-17Rc is greater than 95% as determined by SEC-HPLC.