

Recombinant Cynomolgus TROP-2/TACSTD2 Protein

Catalog No.: RP02682

Sequence Information

Species **Gene ID**
Cynomolgus

Swiss Prot **Tags**
XP_005543292.1 C-His

Synonyms

EGP1; EGP-1; TROP2; GA733-1;
gp50; T16; TACSTD2; TROP-2;
M1S1; TACD2

Product Information

Source **Purification**
HEK293 cells ≥ 95 % as
determined by
Tris-Bis PAGE; ≥
95 % as
determined by
HPLC.

Calculated MW **Observed MW**
28.7 kDa 40-55 kDa

Endotoxin

< 1 EU/μg of the protein by LAL
method.

Formulation

Reconstitution

Centrifuge the tube before
opening. Reconstitute to a
concentration of 0.1-0.5 mg/mL in
sterile distilled water. Avoid vortex
or vigorously pipetting the protein.
For long term storage, it is
recommended to add a carrier
protein or stabilizer (e.g. 0.1% BSA,
5% HSA, 10% FBS or 5%
Trehalose), and aliquot the
reconstituted protein solution to
minimize free-thaw cycles.

Background

Trop-2, also known as epithelial glycoprotein-1 antigen (EGP-1), is a protein that in humans is encoded by the TACSTD2 gene. Mutations of this gene result in gelatinous drop-like corneal dystrophy, an autosomal recessive disorder characterized by severe corneal amyloidosis leading to blindness.

Basic Information

Description

Recombinant Cynomolgus TROP-2/TACSTD2 Protein is expressed from Expi293 with His tag at the C-terminal.; It contains His27-Thr274.

Activity

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Immobilized Cynomolgus TROP-2 at 0.5µg/ml (100µl/Well). Dose response curve for Anti-TROP-2 Antibody, hFc Tag with the EC₅₀ of 10.2ng/ml determined by ELISA.



Recombinant Cynomolgus TROP-2/TACSTD2 Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Cynomolgus TROP-2 is greater than 95% as determined by SEC-HPLC.