

Recombinant Cynomolgus Transferrin R/CD71 Protein

Catalog No.: RP02729

Sequence Information

Species **Gene ID**
Cynomolgus

Swiss Prot **Tags**
XP_005545315.1 N-His

Synonyms
TR; TfR1; CD71; sTfR; T9; p90;
TfRC; TFR

Product Information

Source **Purification**
HEK293 cells ≥ 95 % as
determined by
Tris-Bis PAGE; ≥
95 % as
determined by
HPLC.

Calculated MW **Observed MW**
76.3 kDa 78-80 kDa

Endotoxin
< 1 EU/μg of the protein by LAL
method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

The transferrin receptor (CD71) is known as a receptor for IgA1 on mesangial cells. The transcript and protein expression of glomerular CD71 was determined by real-time polymerase chain reaction and immunohistochemistry. Human mesangial cells (HMCs) were treated with sera from IgAN patients and expression levels of CD71 and inflammatory cytokine markers were compared according to disease status.

Basic Information

Description

Recombinant Cynomolgus Transferrin R/CD71 Protein is expressed from Expi293 with His tag at the N-terminal.; It contains Cys89-Phe760.

Activity

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Recombinant Cynomolgus Transferrin R/CD71 Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Cynomolgus Transferrin R is greater than 95% as determined by SEC-HPLC.